

Effects of Topographical Modification on the Composition and Abundance of Macrofauna in Southern Lagos Lagoon (Ikoyi)

C. A. Brown¹ and E. A. Ajao²

¹Department of Zoology, Marine Biology and Fisheries, University of Lagos, Akoka, Lagos, Nigeria (Email: <ogunwenmo@hotmail.com>)

²Institute of Oceanography and Marine Research, Lagos Island, Lagos, Nigeria

Abstract

The seasonal variation of macrobenthos at Ikoyi was investigated between February 1985 and January 1990. Mean values of surface water temperature, salinity, and percentage mud in sediment and total organic matter were 27.89 °C, 12.22 ‰, 20.36 %, and 5.41 %, respectively. There were no deviations from normal values in a tropical estuarine open lagoon system. Mollusca, Arthropoda, Annelida, Chordata, Nemertina and Chaetognatha were collected. The gastropod mollusc, *Pachymelania aurita*, ranked highest on all but two sampling dates, where *Tympanotonus fuscatus* and *Aloidis trigona* ranked first. Changes in salinity and sediment type were main factors contributing to reduction of species, dominance diversity and change in community structure, which were attributed to habitat modification caused by dredging and sand filling.

Introduction

Lagos lagoon is the largest brackish water body of the southern lagoon system in Nigeria. (Webb, 1958). Although various aspects of the physical properties (Hill & Webb, 1958; Olaniyan, 1969), such as plankton (Olaniyan, 1969; Nwankwo, 1990), epiphytes (Nwankwo & Akinsoji, 1992), finfish (Fagade, 1971, 1979; Fagade & Olaniyan, 1972, 1973, 1974), shell fish (Yoloye, 1974; Yoloye & Adegoke, 1977; Oyekan, 1979; Oyekan & Bolufawi, 1986; Ajao & Fagade, 1990a, 1990b, 1990c, 1990d, 1990e; Brown & Oyekan, 1998), and other fauna (Hill, 1967) have been studied with inputs on pollution (Ekundayo, 1977; Akpata & Ekundayo, 1978; Akpata *et al.*, 1993; Otitolaju, 2002, 2003), as well as mining (Awosika & Dublin-Green, 1994), a census of the present status of the macrobenthic community is necessary due to increased pollution, population explosion

and anthropogenic activities in and around the lagoon.

Longhurst (1958) observed bottom sediments and fauna in the West African continental shelf. Fauna classified by communities belonged to various taxa. Earlier, Smith (1871) described shells from West Africa, hitherto unidentified. Later, Webb (1958) studied the life cycle of *Branchiostoma nigeriense* in the Lagos lagoon. After this there was a dearth of information on benthos in the Lagos lagoon, until Sandison (1966) and Sandison & Hill (1966) observed effects of salinity on macrobenthic fauna in and around the Lagos harbour and adjacent creeks. Yoloye & Adegoke (1977) named a new species of *Neritina* from the Lagos lagoon. The next extensive survey on macrobenthic fauna was not carried out until the next decade when Oyekan (1987) described the benthic macrofaunal communities of the

Lagos lagoon. He described five communities by biotype and biocenosis. More recently, Ajao & Fagade (1990a, 1990b, 1990c) also studied these communities and concluded that the major cause of reduction to extremely low diversities and densities was due to the discharge of various types of untreated wastes into the Lagos lagoon by identified industries. They also recognized a shift in the characteristic fauna in the western industrialized region and eastern unpolluted region. In addition untreated sewage and effluent is discharged into the lagoon (Nwakoro & Odieta, 1997), sand is mined indiscriminately (Awosika & Dublin-Green, 1994), and, in some areas, communities have been completely destroyed by dredging and sand filling activities.

It has, therefore, become imperative to assess the effect of habitat destabilization and modification (caused by dredging activities which evacuate the sediment bottom and sand filling over the mangrove swamp and macrobenthic community), on this localized area in the southern Lagos lagoon, with a view to assessing the problem, proffer solutions and prevent further defaunation.

Materials and methods

Benthic samples were taken in the rainy season (June 1985 and 1986, and July 1987, 1988 and 1989) and dry season (February 1985 and 1986, and January 1988 and 1990) at Ikoyi (Fig. 1), using a 0.1m² Van-Veen grab. No samples were collected from Ikoyi in May 1996 because sand filling and dredging of the surrounding offshore sites made the area too deep to be sampled. Six grab hauls were taken at each station. Benthic fauna were retrieved from five

hauls after Crisp (1971), while the sixth grab haul was used for sediment analysis. Five grab hauls were washed through a 0.5-mm mesh sieve in the field. The retained material was preserved in 5% formalin and kept in labeled jars for further analysis in the laboratory. The sixth grab haul was placed in a labeled polyethylene bag for preservation in the deep freezer in the laboratory for sediment analysis.

Preliminary investigations showed that surface water temperature and bottom water temperature in the Lagos lagoon ranged between 25 °C and 28 °C. Surface water temperatures were, therefore, taken to the nearest whole number, using a mercury-in-glass-thermometer at Ikoyi in the rainy and dry seasons. Water samples were collected with a glass water sampler and stored in clear glass bottles for salinity determination using a temperature compensated hand refractometer (New S-100), in the laboratory. Sediment samples were thawed and dried for 6-8 h at 60-80 °C, to a constant weight. Grain size analysis was carried out as described by Buchanan & Kain (1971). Total organic matter (TOM) was determined using the method described by Oyekan (1981), and the percentage of combustible material in the sediment was estimated as:

$$\frac{\text{Loss of weight on ignition}}{\text{Initial weight before ignition}} \times 100\%$$

The organisms preserved in the field were washed, sorted into taxonomic groups and counted. Faunal density was expressed as species per square metre (m²). Species were ranked and the contributions of the first five numerically dominant species were recorded.




Fig. 1. Map of Lagos lagoon showing Ikoyi sampling station

Margalef's Index (Margalef, 1957), Shannon and Weaver Information Function (Shannon & Weaver, 1963), and Equitability (Lloyd & Ghellardi, 1964) were used to measure diversity in the environment.

Margalef's Index

$$d = \frac{S-1}{\log_e N}$$

d = diversity index, S = number of species,

N = number of individuals, \log_e = natural logarithm.

Shannon and Weaver Information Function

$$H_s = -\sum p_i \log_e p_i$$

S = total number of species; p_i = observed proportion of individuals that belong to the i^{th} species.

Equitability

$$J = \frac{H_s}{\log_2 S}$$

J = equitability measure

H_s = Shannon and Weaver Information Function

S = number of species in sample or community

Results

Physical and chemical properties

Surface water temperature was narrow (26–30 °C) with a mean of 27.89 °C, salinity values were between 0‰ (June and July), and 29‰ (February) with a mean of 12.22‰. Total organic matter was low (< 20.0%) while percentage mud in sediment ranged between 1.3% in February 1986 and 83.2% in February 1985 and had a mean of 5.41% (Table I).

Major animal taxa

The seasonal distribution of phyla at Ikoyi shows a community dominated by molluscs (gastropods and bivalves), which contributed between 77.1% (January) and 100.0% (February 1986) of total fauna collected at

TABLE I

Variation in some physico-chemical parameters during 1985 to 1990

Sampling dates	Feb 1985	June 1985	Feb 1986	June 1986	July 1987	Jan 1988	July 1988	July 1989	Jan 1990	Mean value
Surface water temp. (°C)	27.0	27.0	30.0	27.0	26.0	30.0	28.0	28.0	28.0	27.89
Salinity (‰)	29.0	0.0	27.0	5.0	3.0	21.0	0.0	1.0	24.0	12.22
Mud in sediment (%)	83.2	39.4	1.3	3.7	22.7	19.0	6.9	4.3	2.7	20.36
Total organic matter in sediment (%)	17.8	19.1	1.0	0.5	2.0	4.0	1.0	1.3	2.0	5.41

Ikoyi. (Fig. 2). There was a higher density of gastropods than bivalves. The phylum Annelida contributed between 0.5% (July 1989) and 18.7% (February 1985) of fauna collected. Crustacean arthropods were present and contributed between 0.1% (February 1986) and 23.8% (February 1985). Phylum Nemertina contributed to 4.8% of the sample collected in February 1985. Phylum Chordata contributed 0.1% of sample collected in February 1986 and 0.3% of sample collected in January 1990. The phylum Chaetognatha (*Sagitta* sp.) contributed 0.1% of sample collected in June 1986 (Fig. 2).

Species richness and abundance of individuals

The number of species per m² was highest in February 1985 (26/m²). The density of species collected in January 1990 and July 1988 were 11 and 12, respectively. The density of individuals varied between 10/m² (June 1985) and 60,104/m² (February 1986) (Table 2). A total of 37 different fauna were collected during the sampling period. The gastropod mollusc, *Pachymelania aurita*, was the most abundant species present throughout the sampling period. Densities ranged between 120/m² in February 1985 and 60,056/m² in February 1986.




Fig. 2. Seasonal distribution of phyla at Ikoyi

TABLE 2

Density of macrofauna per m² at Ikoyi for wet and dry seasons between February 1985 and January 1990

	Feb 1985	June 1985	Feb 1986	June 1986	July 1987	Jan 1988	July 1988	July 1989	Jan 1990
NEMERTINA									
Nemertina	28								
ANNELIDA									
<i>Capitella capitata</i>	30								
<i>Glycera convoluta</i>	2						4		
<i>Diopatra neapolitana</i>	10								
<i>Lumbrinereis</i> sp.	12								
<i>Mercierella enigmatica</i>	4								
<i>Nephtys</i> sp.									4
<i>Nereis pelagica</i>				20					
<i>Nereis</i> sp.	12								
<i>Nereis succinea</i>							16	2	46
<i>Phoronis muelleri</i>	2								
<i>Prionospio</i> sp.	2								
<i>Polydora</i> sp.	20								
<i>Syllidae</i>	8								
<i>Tubifex</i> sp.	8								
MOLLUSCA									
<i>Pachymelania aurita</i>	120		60056	1350	1535	1380	564	398	406
<i>Tympanotonus fuscatus</i>	168								
<i>Neritina glabrata</i>			2	68		5	72	16	130
<i>Aloidis trigona</i>	4	4	4			10	6		
<i>Mytilus edulis</i>					5				
<i>Brachyodontes puniceus</i>	4						4		
<i>Cultellus tenuis</i>	6	4							
<i>Diplodonta</i> sp.	2								
<i>Iphigenia truncata</i>	2								4
<i>Crassostrea gasar</i>	4			72		5	20	2	16
ARTHROPODA									
<i>Balanus pallidus</i>	14		32			400			10
<i>Callinectes latimanus</i>	16								
<i>Sersarma huzardi</i>							2		
<i>Clibinarius africanus</i>	32		4			15	8	14	44
<i>Pachygrapsus transversus</i>							2		2
<i>Calappa rubroguttata</i>							4		
<i>Caprella linegris</i>	72								
<i>Corophium insidiosum</i>	4								
<i>Peneaus notialis</i>							12	12	6
Unidentified amphipod	2	2		10					
CHORDATA									
<i>Branchiostoma nigeriense</i>	32								2
CHAETOGNATHA									
<i>Sagitta</i> sp.				2					
Species	26	3	4	7	2	6	12	6	11
Individuals	588	10	60,104	1526	1540	1815	714	444	670

P. aurita ranked first in density among all species collected on all sampling dates except February 1985 when it ranked second to *Tympanotonus fuscatus*. Other high-ranking fauna were *Neritina glabrata*, *Aloidis trigona*, *Crassostrea gazar*, *Nereis succinea*, *Caprella linegris*, *Capitella capitata*, *Nemertina*, *Cutellus tenuis*, unidentified amphipod, *Balanus pallidus*, *Branchiostoma nigeriense*, *Nereis pelagica*, *Mytilus edulis*, *Clibinarius africanus* and *Peneaus notialis*. (Table 3). During the dry season (February 1986), a lagoon visitor, *B. nigeriense* ranked second along side *Balanus pallidus*, while *M. edulis* also a marine species ranked second in July 1987.

Faunal indices

Margalef's Index values ranged between 0.14 in July 1987 and 3.92 in February 1985

(Table 4). Shannon and Weaver Information Function values were highest in February 1985 (2.29) and lowest in February 1986 (0.1) (Table 4). The equitability of individuals ranged from 0.01 (February 1986) to 0.66 (June 1985). (Table 4).

Discussion

The average surface water temperature at Ikoyi was 27.49 °C. This value was higher than that of earlier workers. Fagade (1969) obtained a surface water temperature of 24.8 °C in July 1967 off Ikoyi, while Oyekan (1987) obtained a temperature of 26 °C in October 1974. Salinity was lower in the wet season months (June and July) than the dry season months (January and February). Similar results were obtained from earlier workers (Fagade, 1969; Oyekan, 1987; Brown, 1998).

The *Pachymelania* community was one

TABLE 3
Ranking of macrofauna at Ikoyi between February 1985 and January 1990

	Feb 1985	June 1985	Feb 1986	June 1986	July 1987	Jan 1988	July 1988	July 1989	Jan 1990
<i>Pachymelania aurita</i>	2		1	1	1	1	1	1	1
<i>Tympanotonus fuscatus</i>	1								
<i>Neritina glabrata</i>			4	3		5*	2	2	2
<i>Aloidis trigona</i>		1	3			4			
<i>Crassostrea gazar</i>				2		5*	3	5*	5
<i>Nereis succinea</i>							4	5*	3
<i>Caprella linegris</i>	3								
<i>Capitella capitata</i>	4								
<i>Nemertina</i>	5								
<i>Cutellus tenuis</i>		2							
Unidentified amphipod		3		5					
<i>Balanus pallidus</i>			2*			2			
<i>Branchiostoma nigeriense</i>			2*						
<i>Nereis pelagica</i>				4					
<i>Mytilus edulis</i>					2				
<i>Clibinarius africanus</i>						3		3	4
<i>Peneaus notialis</i>							5	4	

* Same ranking

TABLE 4
Faunal indices at Ikoyi

	Margalef's Index	Shannon & Weaver Information Function	Equitability
Feb. '85	3.92	2.29	0.49
June '85	0.87	1.05	0.66
Feb. '86	0.27	0.01	0.01
June '86	0.83	0.38	0.13
July '87	0.14	0.02	0.02
Jan '88	0.67	0.64	0.25
July '88	1.67	0.99	0.27
July '89	0.82	0.50	0.19
Jan '90	1.54	1.23	0.35

of the five communities described by Oyekan (1987) in the Lagos lagoon; others were Mangrove, Estuarine, Venus and Estuarine rock communities. The *Pachymelania* community inhabited the main part of the Lagos lagoon and consisted of *Pachymelania aurita*, *Aloidis trigona*, *Iphigenia truncata*, *Neritina glabrata*, *Branchiostoma nigeriensis*, *Glycera convoluta*, *Nereis succinea*, *Lumbrinereis* sp., *Diopatra neapolitanea* and crustaceans, *Clibinarius africana* and *Heteropanoe caparti*. *Tympanotonus fuscatus*, *Tagellus angulatus* and *Cultellus tenuis* were also present. Presently in the modified *Pachymelania* community that had reduced in density and diversity from that recorded by Oyekan (1987), Mollusca, Annelida and Arthropoda co-existed alongside Nemertina and Chaetognatha. The latter two were previously absent from this ecosystem. In addition, crabs (*Sersarma huzardi*, *Pachygrapsus transversus* and *Calappa rubroguttata*) were also collected. Their presence in the open lagoon is unusual and suggests a possible change in habitat, which could have been brought

about by sand filling the Mangrove Park at Ikoyi Island. The marine visitor (*Branchiostoma nigeriense*) reported at Ikoyi (Webb, 1958; Brown & Oyekan, 1998) was also collected at Ikoyi during the dry season months. *Balanus pallidus* (a typical mangrove species and fouling organism) also showed abundance during dry season samplings.

One Nemertina, 11 Annelida, three Mollusca and three Arthropoda species collected at Ikoyi in February 1985 were absent

at later sampling dates. The change in community structure from *Pachymelania* community to mangrove community indicates a change in either density or diversity of macrobenthic fauna at Ikoyi. These changes are not independent of physico-chemical factors. Oyekan (1987) observed muddy fine sand at Ikoyi; this later changed to sandy mud (Ajao, 1990d). This could have been attributed to the high diversity observed in February 1985, which dwindled in later sampling dates when the substrate became sandier. Similar reductions in habitat diversity from changes in substrate, current flow and depths (Gordan & Carr, 1978) have also been implicated for shifts and reductions in the number of invertebrate species (Morris *et al.*, 1968).

In addition to substrate type, salinity also determines diversity and density of fauna (Brown & Oyekan, 1998). The salinity at Ikoyi in February 1985 (29.0‰) was the highest collected from all sampling dates. Subsequently, lower salinities (0-27‰) and species diversity (2-12 species/m²) were recorded.

The dredging and sand filling activities at

Ikoyi have pollution implications as they alter bottom conditions over sizeable areas (Ajao & Fagade, 1990e) and destroy mollusc settlement areas (Cole, 1977). These have brought about variations in some physico-chemical parameters and a significant reduction in species, and dominance diversity as well as change in community structure.

Conclusion

The present metamorphosis in community structure provides baseline information on physical, chemical and biological data in southern Lagos lagoon at Ikoyi. This could provide scientific basis for monitoring future ecological changes in the lagoon.

Acknowledgement

The author wishes to acknowledge the assistance rendered by Mr Okunomo and Mr Emubaiye in sample collection during the field trips, Dr J. A. Ajao for academic assistance, and Mr K. Ogunwenmo for reading the manuscript.

References

- Ajao E. A. and Fagade S. O. (1990a). Production and population dynamics of *Pachymelania aurita* Müller. *Arch. Hydrobiol.* **12**(1): 97-109.
- Ajao E. A. and Fagade S. O. (1990b). The ecology of *Capitella capitata* in Lagos lagoon. *Arch. Hydrobiol.* **120**(2): 229-239.
- Ajao E. A. and Fagade S. O. (1990c). The ecology of *Neritina glabrata* in Lagos lagoon, Nigeria. *Arch. Hydrobiol.* **119**(3): 339-350.
- Ajao E. A. and Fagade S. O. (1990d). The distribution and abundance of *Pachymelania aurita* in Lagos lagoon, Nigeria. *Arch. Hydrobiol.* **119**(4): 475-488.
- Ajao E. A. and Fagade S. O. (1990e). A study of the sediments and communities in Lagos lagoon, Nigeria. *J. Oil chem. Poll.* **7**: 85-117.
- Akpata T. V. I., Oyeneke J. A. and Nwankwo D. I. (1993). Impacts of organic pollution on the bacterial, planktonic and benthic populations of Lagos lagoon, Nigeria. *Int. J. Ecol. envir. Sci.* **19**: 73 – 82.
- Akpata T. V. I. and Ekundayo J. A. (1978). Faecal pollution of the Lagos lagoon. *Niger. J. Sci.* **12**(1 & 2): 39 – 53.
- Awosika L. F. and Dublin-Green C. O. (1994). Sand mining in the Lagos and Lekki lagoons and strategies for effective management. *J. Min. Geol.* **30**(11): 137-139.
- Brown C. A. (1998). Distribution and population dynamics of an estuarine population of *Aloidis trigona* Hinds. (Bivalvia). *Acta Hydrobiol.* **40**: 227 – 237.
- Brown C. A. and Oyeneke J. A. (1998). Temporal variability in the structure of benthic macrofauna communities of the Lagos lagoon and harbour, Nigeria. *Pol. Arch. Hydrobiol.* **45**(1): 45-54.
- Buchanan J. B. and Kain J. B. (1971). Measurement of physical and chemical environment. In *Methods for the study of marine benthos* (N. A. Holmes and A.D. McIntyre, ed.), pp. 30-58. Blackwell Scientific Publication.
- Cole H. A. (1977). Sand and gravel extraction. *Marine Pollution Bull.* **8**. 193 pp.
- Crisp D. J. (1971). Energy flow measurements. In *Methods for the study of marine benthos*. (N. A. Holme and A. D. McIntyre, ed.), pp. 197 – 279. Blackwell Scientific Publications.
- Ekundayo J. A. (1977). Environmental consequences of pollution of the Lagos lagoon. *Bull. Sci. Ass. Niger.* **3**(2): 290-299.
- Fagade S. O. (1969). *Studies on biology of some fishes of Lagos lagoon*. (PhD Thesis.) University of Lagos, Nigeria. 358 pp.
- Fagade S. O. (1971). The food and feeding habits of *Tilapia* species in Lagos lagoon. *J. Fish. Biol.* **3**: 151-156.
- Fagade S. O. (1979). Observations on the biology of two species of *Tilapia* from the Lagos lagoon, Nigeria. *Bull. D'IFAN* **41** (1): 629-654.
- Fagade S. O. and Olaniyan C. I. O. (1972). The

- biology of the West African Shad, *Ethmalosa ufbriata* (Bowdich) in the Lagos lagoon, Nigeria. *J. Fish. Biol.* **4**: 519-534.
- Fagade S. O. and Olaniyan C. I. O.** (1973). The food and feeding interrelationship of the fishes in the Lagos lagoon. *J. Fish. Biol.* **5**: 205-225.
- Fagade S. O. and Olaniyan C. I. O.** (1974). Seasonal distribution of the fish fauna of the Lagos lagoon. *Bull. d'IFAN* **36**(1): 244-252.
- Gordan O. T. and Carr J. R.** (1978). Habitat structure and stream fish communities. *Ecology* **59**: 507-515.
- Hill M. B.** (1967). The life cycles and salinity tolerance of the serpulids *Mercierella enigmatica* (Fauvel) and *Hydroides unciata* (Philipi) at Lagos, Nigeria. *J. Anim. Ecol.* **36**: 306-321.
- Hill M. N. and Webb J. E.** (1985). The ecology of Lagos lagoon II. The topography and physical features of Lagos harbour and Lagos lagoon. *Phil. Trans. R. Soc.* **241**: 307-419.
- Lloyd M. and Ghellardi R. J.** (1964) A table for calculating the equitability components for species diversity. *J. Anim. Ecol.* **33**: 217-225.
- Longhurst A. R.** (1958). An ecological survey of the west African Marine Benthos. *Fishery Publs Lond.* **11**: 1-102.
- Margalef R.** (1957). La teoria de la informacion enecologia. *Mems. R. Acad.* **32**: 273-449.
- Morris L. A., Longermier R. N., Russel T. R. and Witt A. Jr.** (1968). Effects of the mainstream impoundments and channelisation upon the limnology of the Missouri River, Nebraska. *Trans. Am. Fish. Soc.* **97**(4): 380-388.
- Nwankwo D. I.** (1990). Distribution and seasonal variation of dinoflagellates in Lagos lagoon. *Niger. J. Bot.* **3**(53): 197-208.
- Nwankwo D. I. and Akinsoji A.** (1992). Epiphyte community on water hyacinth *Eichhornia crassipes* (Mart) Solms in coastal waters of southwestern Nigeria. *Arch. Hydrobiol.* **124** (4): 501-511.
- Nwaokoro R. C. and Odieta W. O.** (1997). Baseline survey of industries, industrial wastewaters and waterbodies receiving industrial wastewaters in Lagos State. *World Bank funded project for Ministry of Environmental & Physical Planning. Ikeja.* 77 pp.
- Olaniyan C. I. O.** (1969). The seasonal variation in the hydrology and total plankton of the lagoons of southwest Nigeria. *Niger. J. Sci.* **3**(2): 101-119.
- Otitoloju A. A.** (2002). Evaluation of joint action toxicity of binary mixtures of heavy metals against the mangrove periwinkle *Tympanotonus fuscatus* var. *radula* (L.). *Ecotoxicol. envir. Safety.* **53**: 404-415.
- Otitoloju A. A.** (2003). Relevance of joint action toxicity evaluations in setting realistic environmental safe limits of heavy metals. *J. envir. Mgmt* **67**: 121-128.
- Oyenekan J. A.** (1979). The ecology of the genus *Pachymelania* in Lagos lagoon. *Arch. Hydrobiol.* **86**(4): 515-522.
- Oyenekan J. A.** (1981). *Community structure and production of benthic macrofauna of Southampton water.* (PhD Thesis.). University of Southampton. 351 pp.
- Oyenekan J. A.** (1987). Benthic macrofaunal communities of Lagos lagoon, Nigeria. *Niger. J. Sci.* **21**: 45-51.
- Oyenekan J. A. and Bolufawi J. E.** (1986). The ecology of *Iphigenia truncata* in Lagos lagoon. *Arch. Hydrobiol.* **106**(4): 559-566.
- Sandison E. E.** (1966). The effect of salinity fluctuations on the life of *Balanus pallidus* Strusburi (Darwin) in Lagos harbour, Nigeria. *J. Anim. Ecol.* **35**: 363 - 378.
- Sandison E. E. and Hill M. B.** (1966). The distribution of *Balanus pallidus* Strutburi Darwin, *Gryphea gazer* Adanson, Dautzenbury, *Mercierella enigmatica* Fauvel and *Hydroides uncinata* (Phillippi) in relation to salinity in Lagos harbour and adjacent creeks. *J. Anim. Ecol.* **35**: 235 - 258.
- Shannon C. E. and Weaver W.** (1963). *The Mathematical Theory of Communication.* University of Illinois Press. Urbana. 125 pp.
- Smith E. A.** (1871). A list of species of shells

- from West Africa, with descriptions of those hitherto described. *Bull. Brit. Mus. Nat. Hist. Zool.* 727-739.
- Webb J. E.** (1958). The ecology of Lagos lagoon. I. The lagoon of the Guinea Coast. *Phil. Trans. R. Soc. Lond.* 683(241): 307-318.
- Williams N. V.** (1962). The seasonal distribution of the teleost fish in Lagos harbour, creek and lagoon in relation to salt tolerance (MSc Thesis). University College, North Wales. Bangor. pp. 80-90.
- Yoloye V. L.** (1974). The sexual phase of the West African cockle *Anadara senilis*. *Proc. malac. Soc. Lond.* 41: 25-28.
- Yoloye V. and Adegoke S.** (1977). A new species of *Neritina*. (Archaeogastropoda, Neritidae) from the Lagos lagoon. *Malacologia* 16(1): 303-309.